



Pharmacy Prior Authorization Botulinum Toxins – Clinical Guideline

Botox (onabotulinumtoxinA)
Dysport (abobotulinumtoxinA)

Myobloc (rimabotulinumtoxinB)
Xeomin (incobotulinumtoxinA)

Prior Authorization Guidelines for All Indications:

Botox, Myobloc, Dysport, and Xeomin must be prescribed by an appropriate specialist based on indication, and meet the following criteria:

- **Migraine Prophylaxis (*Botox*):**
 - Prevention of chronic migraine
 - Fifteen or more days per month with headaches lasting 4 hours a day or longer
 - Member had inadequate response to, or intolerable side effects, with at least two medications from two classes of migraine headache prophylaxis, for at least two months (60 days):
 - Beta-blocker: propranolol, metoprolol, timolol, atenolol, nadolol
 - Anticonvulsant: valproic acid or divalproex, topiramate
 - Antidepressants: amitriptyline, venlafaxine, duloxetine
 - Member is at least 18 years of age
 - Medication will not be used concurrently with calcitonin gene-related peptide (CGRP) receptor antagonists
- **Chronic Limb Spasticity (*Botox, Xeomin, Dysport*):**
 - Spasticity may be due to an injury to the brain or spinal cord, or along with a neurological disorder
 - For example, stroke, traumatic brain injury, multiple sclerosis, spinal cord injury, cerebral palsy
 - Member had previous trial and failure with baclofen
 - Trial of physical therapy and/or occupational therapy
 - Botox and Dysport: Member is at least 2 years of age
 - Xeomin:
 - Upper limb spasticity, excluding spasticity caused by cerebral palsy: Member is at least 2 years of age
 - All other limb spasticity, member is at least 18 years of age
- **Focal Spasticity or Equinus Gait due to Cerebral Palsy (*Botox, Dysport*):**
 - Member is 2 years of age or older
- **Severe primary axillary hyperhidrosis - excessive underarm sweating (*Botox, Dysport*):**



Pharmacy Prior Authorization Botulinum Toxins – Clinical Guideline

- There is focal, visible, excessive sweating for at least 6 months without apparent cause and two of the following:
 - Interferes with daily activities
 - Bilateral and relatively symmetric
 - Onset before 25 years of age
 - Focal sweating stops during sleep
 - Family history of idiopathic hyperhidrosis
 - At least one episode per week
- Failure of topical aluminum chloride (hexahydrate)
- Member is at least 18 years of age
- **Neurogenic bladder (*Botox*):**
 - Diagnosis of urinary incontinence due to detrusor overactivity associated with neurologic condition
 - Trial of behavioral therapy (for example, bladder training, bladder control strategies, pelvic floor muscle training, fluid management) for at least 8-12 weeks
 - Trial and failure or intolerance with two formulary urinary anticholinergics
 - For example, oxybutynin, trospium, tolterodine, solifenacin
 - Member is at least 5 years of age
- **Overactive bladder (*Botox*):**
 - Trial of behavioral therapy (bladder training, bladder control strategies, pelvic floor muscle training, fluid management) for at least 8-12 weeks
 - Trial and failure or intolerance with two formulary urinary anticholinergics
 - For example, oxybutynin, trospium, tolterodine, solifenacin
 - Member is at least 18 years of age
- **Esophageal Achalasia (*Botox*):**
 - Member meets ONE of the following:
 - Member remains symptomatic despite surgical myotomy or pneumatic dilation
 - Member is at high surgical risk or unwilling to undergo surgical myotomy or pneumatic dilation
 - Member is at least 18 years of age



Pharmacy Prior Authorization Botulinum Toxins – Clinical Guideline

- **Chronic anal fissures (*Botox*):**
 - Trial and failure of nitroglycerin ointment 0.4% (Rectiv) AND either bulk fiber supplements, stool softeners, or sitz baths for at least two months
 - Endoscopy has been completed to rule out Crohn's disease
 - Member is at least 18 years of age
- **Chronic sialorrhea - excessive drooling (*Botox, Myobloc, Xeomin*):**
 - Trial and failure of anticholinergic such as glycopyrrolate (pediatric use ages 3-16 years) or benztropine (adults)
 - Botox: Member is at least 21 months of age
 - Xeomin: Member is at least 2 years of age
 - Myobloc: Member is at least 18 years of age

Botulinum toxins may also be approved if medically necessary for treatment of the following indications having limited treatment options:

- Botox for cervical dystonia (spasmodic torticollis):
 - Member at least 16 years of age
- Dysport, Myobloc, Xeomin for cervical dystonia:
 - Member is at least 18 years of age
- Botox for blepharospasm associated with dystonia, including benign essential blepharospasm or VII nerve disorders:
 - Member is at least 12 years of age
- Xeomin for blepharospasm:
 - Member is at least 18 years of age and previously treated with onabotulinumtoxinA (Botox)
- Dysport for blepharospasm:
 - Member is at least 18 years of age and previously treated with onabotulinumtoxinA (Botox) and incobotulinumtoxinA (Xeomin)
- Botox for strabismus:
 - Member is at least 12 years of age
- Botox for hemifacial spasm:
 - Member is at least 18 years of age



Pharmacy Prior Authorization Botulinum Toxins – Clinical Guideline

Initial Approval:

- 6 months - Treatment is given every 12 weeks

Renewal Approval:

- 1 year - Treatment is given once every 12 weeks
- Botox:
 - Does not exceed a cumulative dose of 400 units every 90 days for adults
 - Does not exceed the lower of 10 units per kg or 340 units every 90 days for pediatric patients

Additional Information:

- If members do not respond to a course of treatment (usually lasts for 12 weeks), treatment should be discontinued.
- Continuing treatment with botulinum toxin injection for ongoing prevention of chronic migraine headaches is considered medically necessary when:
 - Migraine headache frequency was reduced by at least 7 days per month (when compared to pre-treatment average) by the end of the initial trial; OR
 - Migraine headache duration was reduced by at least 100 total hours per month (when compared to the pre-treatment average) by the end of the initial trial

References:

1. Botox® (onabotulinumtoxinA) [package insert]. Allergan, Inc, Madison, NJ; Revised July 2021. <https://media.allergan.com/actavis/actavis/media/allergan-pdf-documents/product-prescribing/20190620-BOTOX-100-and-200-Units-v3-0USPI1145-v2-0MG1145.pdf>. Accessed November 11, 2021.
2. Dysport® (abobotulinumtoxinA) [package insert]. Wrexham, UK: Ipsen Biopharm, Ltd.; Revised July 2020. https://www.ipsen.com/websites/Ipsen_Online/wp-content/uploads/2020/07/10002305/DYS-US-004998_Dysport-PI-July-2020.pdf. Accessed November 11, 2021.
3. Xeomin® (incobotulinumtoxinA) [package insert]. Merz Pharmaceuticals, LLC, Raleigh, NC; Revised August 2021. <https://dailymed.nlm.nih.gov/dailymed/fda/fdaDrugXsl.cfm?setid=ccdc3aae-6e2d-4cd0-a51c-8375bf9458&type=display>. Accessed November 11, 2021.
4. Myobloc® (rimabotulinumtoxinB) [package insert]. South San Francisco, CA: Solstice Neurosciences, Inc.; Revised March 2021. https://myobloc.com/files/MYOBLOC_PI.pdf. Accessed November 11, 2021.
5. Simpson DM, Hallet M, Ashman EJ, et al. Practice guideline update summary: botulinum neurotoxin for the treatment of blepharospasm, cervical dystonia, adult spasticity, and headache: Report of the guideline development subcommittee of the American Academy of Neurology. *Neurology*. 2016;86:1818-1826.
6. Yelnik AP, Simon O, Bensmail D, et al. Drug treatments for spasticity. *Annals Phys Rehab Med*. 2009;52(10):746-756.
7. Smith C, Pariser D. Primary Focal Hyperhidrosis. Waltham, MA. UpToDate. Last Modified January 2, 2020. <https://www.uptodate.com/contents/primary-focal-hyperhidrosis>. Accessed August 19, 2020.
8. Evers S, Afra J, Frese A, Goadsby PJ, et al. EFNS guideline on the drug treatment of migraine – revised report of an EFNS task force. *Eur J Neurol*. 2009;16: 968–981.
9. S.D. Silberstein, S. Holland, F. Freitag, et al. Evidence-based guideline update: Pharmacologic treatment for episodic migraine prevention in adults. *Neurology* 2012;78;1337-1345.



Pharmacy Prior Authorization Botulinum Toxins – Clinical Guideline

10. Estemalik E, Tepper S. Preventive treatment in migraine and the new US guidelines. *Neuropsychiatric Disease and Treat.* 2013;9:709–720.
11. Olsson JE, Behrin HC, Forsman B, et al. Metoprolol and propranolol in migraine prophylaxis: a double-blind multicenter study. *Acta Neurol Scand.* 1984;70(3):160-168.
12. Dodick DW, et al. PREEMPT Chronic Migraine Study Group. OnabotulinumtoxinA for treatment of chronic migraine: pooled results from the double-blind, randomized, placebo-controlled phases of the PREEMPT clinical program. *Headache.* 2010;50(6):921.
13. Mahajan ST. Use of Botulinum Toxin for Treatment of Non-neurogenic Lower Urinary Tract Conditions. Waltham, MA. UpToDate. Last Modified: February 5, 2020. <https://www.uptodate.com/contents/use-of-botulinum-toxin-for-treatment-of-non-neurogenic-lower-urinary-tract-conditions>. Accessed August 19, 2020.
14. Ates F, Vaezi MF. The pathogenesis and management of achalasia: current status and future directions. *Gut Liver.* 2015;9(4):449-463.
15. Vaezi MF, Pandolfino JE, Vela MF. ACG clinical guideline: Diagnosis and management of achalasia. *American Journal of Gastroenterology.* 2013; 108:1238-1249.
16. Wald A, Bharucha AE, Cosman BC, Whitehead WE. ACG clinical guideline: management of benign anorectal disorders. *Am J Gastroenterol.* 2014;109:1141-1157.
17. Bleday R. Anal Fissure: Medical Management. Waltham, MA. UpToDate. Last Modified February 24, 2020. <https://www.uptodate.com/contents/anal-fissure-medical-management>. Accessed August 19, 2020.
18. Reddihough D, Erasmus CE, Johnson H, McKellar GMW, Jongerius PH. Botulinum toxin assessment, intervention and aftercare for pediatric and adult drooling: international consensus statement. *Eur J Neurol.* 2010;17(Suppl 2):109-121.
19. Jongerius PH, van den Hoogen FJA, van Limbeek J, Gabreels FJ, van Hulst K, Rotteveel JJ. Effect of Botulinum Toxin in the Treatment of Drooling: A Controlled Clinical Trial. *Pediatrics.* 2004; 114(3): 620 -627.
20. Reddihough D, Erasmus CE, Johnson H, McKellar GMW, Jongerius PH. Botulinum toxin assessment, intervention and aftercare for pediatric and adult drooling: international consensus statement. *Eur J Neurol.* 2010;17(Suppl 2):109-121.
21. Walshe M, Smith M, Pennington L. Interventions for drooling in children with cerebral palsy. *Cochrane DB Syst Rev.* 2012, Issue 11. Art. No.: CD008624. DOI: 10.1002/14651858.CD008624.pub3.
22. Jongerius PH, van den Hoogen FJA, van Limbeek J, Gabreels FJ, van Hulst K, Rotteveel JJ. Effect of Botulinum Toxin in the Treatment of Drooling: A Controlled Clinical Trial. *Pediatrics.* 2004; 114(3): 620 -627.
23. Costa J, et al. Botulinum toxin type A therapy for blepharospasm (Review). *COCHRANE DB SYST REV.* 2004, Issue 2. Art.No.: CD004900. DOI: 10.1002/14651858.CD004900.pub2.
24. Patterson MC. Management and prognosis of cerebral palsy. UpToDate. <http://www.uptodate.com>. Updated April 8, 2016. Accessed August 19, 2020.
25. Pavone V, Testa G, Restivo DA, et al. Botulinum toxin treatment for limb spasticity in childhood cerebral palsy. *Front Pharmacol.* 2016;7:29.
26. Koman LA, Brashear A, Rosenfeld S, et al: Botulinum toxin type a neuromuscular blockade in the treatment of equinus foot deformity in cerebral palsy: a multicenter, open-label clinical trial. *Pediatrics* 2001; 108(5):1062-1071.
27. AbobotulinumtoxinA for equinus foot deformity in cerebral palsy: a randomized controlled trial. 2016;137(2): e20152830.
28. Delgado MR, Hirtz D, Aisen M, et al. Practice parameter: pharmacologic treatment of spasticity in children and adolescents with cerebral palsy (an evidence-based review): report of the Quality Standards Subcommittee of the American Academy of Neurology and the Practice Committee of the Child Neurology Society. *Neuro.* 2010;74(4):336.
29. Hoare BJ, Wallen MA, Imms C, Villanueva E, Rawicki HB, Carey L. Botulinum toxin A as an adjunct to treatment in the management of the upper limb in children with spastic cerebral palsy (update). *Cochrane Database Syst Rev.* 2010;(1): CD003469. doi: 10.1002/14651858.CD003469.pub4.
30. Barkoudah E, Glader L. Cerebral palsy: Treatment of spasticity, dystonia, and associated orthopedic issues; Waltham, MA. UpToDate. October 2021; https://www.uptodate.com/contents/cerebral-palsy-treatment-of-spasticity-dystonia-and-associated-orthopedic-issues?search=chronic%20limb%20spasticity%20treatment&source=search_result&selectedTitle=1~150&usage_type=default&display_rank=1#H2514948445. Accessed November 11, 2021.
31. Smith C, Pariser D, Primary focal hyperhidrosis; Waltham, MA. UpToDate. October 2021; https://www.uptodate.com/contents/primary-focal-hyperhidrosis?search=Severe%20primary%20axillary%20hyperhidrosis%20&source=search_result&selectedTitle=2~150&usage_type=default&display_rank=2#H197353549. Accessed November 11, 2021.



Pharmacy Prior Authorization Botulinum Toxins – Clinical Guideline

32. Cephalalgia. 2002 Sep;22(7):491-512. doi: 10.1046/j.1468-2982.2002.00386.x. Migraine: preventive treatment; S D Silberstein 1, P J Goadsby
33. Schwedt T, Garza I; Preventive treatment of episodic migraine in adults; Waltham, MA. UpToDate October 2021; https://www.uptodate.com/contents/preventive-treatment-of-episodic-migraine-in-adults?search=botulinum%20toxins%20migraine%20prophylaxis&source=search_result&selectedTitle=1~150&usage_type=default&display_rank=1. Accessed November 11, 2021.
34. Fehlings D, Rang M, Glazier J, et al: An evaluation of botulinum-A toxin injections to improve upper extremity function in children with hemiplegic cerebral palsy. *The Journal of Pediatrics*. 2000; 137:331-337.
35. Gold Standard. (2016, Jan 26). Botox. Tampa, Florida, USA. Retrieved August 16, 2017, from <http://www.clinicalpharmacology-ip.com/Forms/Monograph/monograph.aspx?cpnum=1219&sec=monindi&t=0>
36. Gormley, EA, Lightner, DJ, Burgio, KL, et al. Diagnosis and Treatment of Overactive Bladder (non-Neurogenic) in Adults:AUA/SUFU guideline. *The Journal of Urology*, 2012: 188 (6), 2455-2463.
37. Abrams, GM, Wakasa, M. Chronic Complications of Spinal Cord Injury and Disease. Waltham, MA. UpToDate. Last Modified December 6, 2018. <https://www.uptodate.com/contents/chronic-complications-of-spinal-cord-injury-and-disease>. Accessed August 14, 2019.
38. Egevad. G. Petkova, Y, Villholm, O. Sialorrhea in Patients with Parkinson's Disease: Safety and Administration of Botulinum Neurotoxin. *Journal of Parkinson's Disease*. 2014 :4 (3): 321-326.
39. American Academy of Neurology. <https://www.aan.com/Guidelines/home/GetGuidelineContent/377>. Accessed August 2, 2018.
40. National Institute of Neurological Disorders and Stroke. Spasticity information page. <https://www.ninds.nih.gov/Disorders/All-Disorders/Spasticity-Information-Page>. Accessed August 19, 2020.